

VitaLink Grow Schedules

ADDITIVE RANGE



* VitaLink Fulvic is also recommended to be used as a foliar application. Dilute 10 ml/L and apply once a week.

** When using VitaLink Coir, please use VitaLink Buddy at 5ml/10L.

Note – When growing conditions are hot (<27oC) and dry (>40%RH), maintain the conductivity at the lower end of the recommendation above.



BIOPLUS



BIOPAC



FULVIC



FOLIAR



BUDDY



FLUSH

Propagation Stage 1

Pre-soaking plugs

2 ml/L

50 ml/L

-

-

-

-

Propagation Stage 2

Rooted cuttings/seedlings

2 ml/L

50 ml/L

-

-

-

-

Vegetative Stage 1 - Duration 1 to 2 weeks

Establishment period

1 ml/10 L

15 ml/50 L

2 ml/L*

100 ml/L

-

-

Vegetative Stage 2 - Duration 1 to 2 weeks

Aggressive growth

1 ml/10 L

15 ml/50 L

2 ml/L*

100 ml/L

-

-

Reproductive Stage 1 - Duration 1 to 2 weeks

First signs of flowers

1 ml/10 L

15 ml/50 L

2 ml/L*

100 ml/L

-

-

Reproductive Stage 2 - Duration 2 weeks

Develop small flowers/fruit

1 ml/10 L

15 ml/50 L

2 ml/L*

100 ml/L

-

-

Reproductive Stage 3 - Duration 2-3 weeks

Expansion in size of flowers/fruit

1 ml/10 L

15 ml/50 L

2 ml/L*

100 ml/L

1 ml/L**

-

Reproductive stage 4 – Duration 1-2 weeks

Flowers/Fruit ripen

1 ml/10 L

15 ml/50 L

2 ml/L*

100 ml/L

1 ml/L**

4 ml/L